

MICROFICHE N° 826 A TIR DE
L'UNITE DOCUMENTAIRE
N

جديدة منجزة حسب الوثيقة
رقم :

93

0095

ROYAUME DU MAROC


المملكة المغربية

المركز المغربي للدراسات والبحوث



CENTRE NATIONAL DE DOCUMENTATION


SERVICE DE REPROGRAPHIE
ET IMPRIMERIE

B.P. 826 RABAT



مصلحة الطباعة والتصوير
ص.ب 826 الرباط





WATER USE EFFICIENCY, GROWTH AND DEVELOPMENT RESPONSE OF MAIZE TO PLANT POPULATIONS IN SEMI-ARID MOROCCO

M. KARROU*, H. MINOR and L. CROY*

INTRODUCTION

In semi-arid areas of the world, soil water deficits and excessively high temperatures are probably the most common yield - limiting factors in crops. Among these crops, maize yield is limited most by drought that occurs during its reproductive period. To improve yield, many scientists are seeking means of reducing the drought effect and hence making corn more water use efficient. Controlling plant population is one of the important practices to match water use to anticipated availability of soil moisture (Waldren, 1983). Timmons et al. (1966) demonstrated that when precipitation was below average from planting through silking, different stand depleted available soil moisture to near the wilting point at later stage and evapotranspiration remained about the same for each population so the efficiency of water use varied with yield. However, no definite relationship was found by Bababola and Oputa (1981) between water

* Cereal Agronomists, Aridoculture center, INRA/MIAC, B.P. 290, Sétif, Morocco

** Associate Professor, Dep. of Agronomy, Univ. of Missouri, Columbia (U.S.A.)

use efficiency and population necessary for optimum grain yields since optimum corn stands were related to growing seasons.

Increasing plant population from about 30,000 to 70,000 plants/ha in adverse soil water seasons decreased grain yields of corn (Bababdola and Oputa, 1981) and reduced effective filling period duration by 2.5 days at high plant density (Ponelcic and Egli, 1979). Claassen and Shaw (1970) postulated that reductions in yields were due to reductions in kernel numbers when stress occurred before or during silking and pollination, and to reduction in kernel weights when stress occurred after silking. However, Harden et al., (1982) concluded that moisture stress within two weeks after silking reduced the number of kernels /plant by about 15% with little influence on final kernel weight. Later stress cycles had only minor effects on kernel number.

In semi-arid Morocco, one of the techniques used by farmers to reduce plant competition for water during grain filling and to feed their livestock is the thinning of less vigorous plants at tasseling. The objective of this study was to determine if plant population level obtained either by seeding rate or by thinning plants at tasseling (practice of Moroccan farmers) can reduce the competition among plants and conserve water in the soil for use during the reproductive and grain filling periods when moisture deficiency usually occurs.

MATERIALS AND METHODS

Experiments were conducted at Sidi El Aydi and Jemaa Shaim experiment stations, located in Chaouia and Abda plains (Morocco), respectively. Soils on which experiments were located are alkaline vertisols (pH7.5-8.0). Mean annual precipitation was 388 and 318 mm at Sidi El Aydi and Jemaa Shaim, respectively. The hybrid used was a moroccan double cross DRA 400 planted Feb 1, 1985 in Sidi El Aydi and 7 Feb 12, 1985 in Jemaa Shaim. Treatments consisted of four expected plant populations or densities : low -20,000 (d1); medium -40,000 (d2); high -80,000 plants / ha thinned at tasseling to 40,000 plants / ha (d4). Data collected were, total dry matter, grain yield, yield components, midday leaf water potential, leaf diffusive resistance, leaf transpiration, and soil moisture. Seeds were planted at 3 seeds / hill and then plants were thinned to 1 per hill at the 4 leaf stage. Row spacing was 75 cm. The experimental design was a randomized complete block design with 4 replications.

RESULTS AND DISCUSSION

Water status

Total water used from emergence to physiological maturity was not affected by plant population (Table 1). Increasing plant population apparently hastened soil water depletion. Leaf transpiration (Fig.1) decreased earlier for the high plant population than for the lower ones. Leaf water potential (Fig.2) declined over time as the season progressed in 1985. Curves were parallel in the case of the low and medium plant populations. For the high plant population, early the curve was parallel to the other curves, then the rate of decrease was very fast after day 100 (29 May, silking stage). The lowest leaf water potentials were -1.9, -2.2 and -3.3 MPa for d1, d2 and d3, respectively, when the plants were at the dent stage. The critical value of the leaf water potential for which leaves could not recover at night was around -2.2 MPa in this study, but was never reached by the low plant population.

Table I : Total water used (cm) from emergence to physiological maturity at Sidi El Aydi (SEA) and Jemaa Shaim (JS) in 1985 and 1986.

Treatment Plants/ha	Cropping Season and Location			
	SEA 85	JS 85	SEA 86	JS 86
d1 : 20.000	14.70	10.71	10.14	13.88
d2 : 40.000	15.36	11.27	10.07	15.24
d3 : 80.000	15.26	12.64	11.38	13.88
d4 : 80.000 + Thinning	14.46	11.14	10.07	13.87
LSD (0.05)	n.s.	n.s.	---	n.s.
LSD (0.01)	n.s.	n.s.	---	---
CV (%)	8.90	19.37	11.30	12.41




Fig. 1 : Leaf transpiration versus time at Sidi El Aydi.




Fig. 2 : Leaf water potential versus time at Sidi El Aydi

The low plant population (d1) did not induce variation in total leaf diffusive resistance until the end of the growing season, where a small increase was measured (Fig.3). Leaf diffusive resistance remained constant until silking stage (100 days after emergence) for the other two plant populations (d2 and d3) and then increased rapidly. This increase was faster for d3 than d2 and corresponded to the rapid decrease of leaf water potential.




Fig. 3 : Leaf diffusive Resistance versus time at Sidi El Aydi

To gain an understanding of the relationship of stomata closure and leaf water potential, leaf diffusive resistance was plotted against leaf water potential (Fig. 4). The lack of increase of leaf diffusive resistance when leaf water potential decreased from -0.3 to -1.6 MPa can be explained by drought that occurred during March and April of 1985. This drought probably hardened plants and made them less sensitive to leaf water potential variation. Nevertheless, values of -1.6 to -2.0 MPa and 4 s cm⁻¹ correspond to the beginning of stomatal closure. These values are similar to those reported in the literature (Boyer, 1970)

Yields

Grain yield and its components are presented in Tables 2, 3 and 4, respectively. Numbers of kernels / ear decreased with the increase of plant




Table II : Grain yield at Sidi El Aydi (SEA) and Jemaa Shaim (JS) in 1985 and 1986

Treatment Plants/ha	Cropping Season and Location			
	SEA 85	JS 85	SEA 86	JS 86
			Qx/ha	
d1 : 20.000	13.98	8.85	3.85	4.15
d2 : 40.000	16.98	12.46	1.97	5.91
d3 : 80.000	16.78	14.16	1.48	5.27
d4 : 80.000 + Thinning	19.64	11.73	1.92	4.42
LSD (0.05)	2.67	3.10	0.83	n.s.
LSD (0.01)	3.83	4.46	1.26	n.s.
CV (%)	9.96	16.44	18.14	17.18

Table III : Number of kernels/ear at Sidi El Aydi (SEA) and Jemaa Shaim (JS) in 1985 and 1986.

Treatment	Cropping Season and Location			
	SEA 85	JS 85	SEA 86	JS 86
			# Kernels / Ear	
d1 : 20.000	450	454	312	338
d2 : 40.000	415	435	304	377
d3 : 80.000	365	405	222	302
d4 : 80.000 + Thinning	413	455	254	318
LSD (0.05)	32	n.s.	n.s.	n.s.
LSD (0.01)	46	n.s.	--	--
CV (%)	5	9	15	14

Table IV : Seed weight at Sidi El Aydi (SEA) and Jemaa Shaim (JS) in 1985
and 1986

Treatment	Cropping Season and Location			
	SEA 85	JS 85	SEA 86	JS 86
	mg / Kernel			
d1 : 20.000	21.4	24.4	13.1	13.7
d2 : 40.000	18.2	20.9	12.4	13.4
d3 : 80.000	15.5	19.3	12.8	12.5
d4 : 80.000 + Thinning	17.1	19.6	12.7	12.5
LSD (0.05)	2.0	3.2	n.s.	0.8
LSD (0.01)	2.8	4.5	n.s.	1.2
CV (%)	6.8	9.4	6.5	3.1

Table V : Aboveground yield at Sidi El Aydi (SEA) and Jemaa Shaim (JS) in 1985 and 1986.

Treatment	Cropping Season and Location			
	SEA 85	JS 85	SEA 86	JS 86
	Qx / ha			
d1 : 20.000	32.04	16.60	21.11	20.83
d2 : 40.000	41.31	21.96	24.55	36.61
d3 : 80.000	45.98	27.39	22.33	45.50
d4 : 80.000 + Thinning	45.07	25.49	21.45	29.06
LSD (0.05)	9.90	--	n.s.	14.60
LSD (0.01)	15.00	--	n.s.	22.12
CV (%)	12.01	--	9.24	16.51

population. However, these decreases remained statistically not significant in 1985 at Jemaa Shaim and in 1986 at both sites. Nevertheless, seed weight decreased roughly with the increase of plant population in all conditions. Meanwhile, the differences were highly significant in 1985 at Sidi el Aydi. The reason was that 1986 was too dry at the end of the cycle and affected all plant populations. However, in 1985 the cropping season was relatively wet and only the highest plant population was affected. Leaf transpiration was very low in the case of the high plant population after silking (Fig. 1). In 1985, the highest yields obtained by the medium and high plant population were not statistically different. However, high seed weight and number of kernels/ear obtained with the low plant population were unable to compensate for the lower plant population. So the yields remained low. In 1986, these components of yield were too low; and consequently, yields were also very low.

Thinning of plants at tasseling did not improve yields and its components during a dry year (1986) because most of the water stored in the soil was depleted at thinning time. In a relatively wet cropping season (1985), thinning improved the yield and its components, especially at Sidi El Aydi.

Aboveground yield and harvest indices are presented in Tables 5 and 6. These parameters increased and decreased respectively with the increase of plant population in the two locations during the two-year experiment. Data on harvest index show that more dry matter is accumulated in the grain as plant population becomes smaller. However, the harvest indices obtained in 1986 are very low because of the rapid growth before tasseling that induced a rapid depletion of soil moisture. Consequently, the reproductive period was very dry and grain yield was very low.

Water Use Efficiency

Water use efficiency is the ratio of dry matter produced to the amount of water used. Because the amount of water used by different plant populations was similar, the efficiency of water use varied with yield. In fact, the low and medium (d2) plant populations used water more efficiently during dry and wet cropping seasons, respectively (Table 7).

From this study we can conclude that optimum plant population is 20,000 plants/ha and 40,000 plants/ha in a dry and wet year, respectively. However, thinning should be done earlier during a dry year.

Table VI : Harvest indices at Sidi El Aydi (SEA) and Jemaa Shaim (JS) in 1985 and 1986.

Treatment	Cropping Season and Location			
	SEA 85	JS 85	SEA 86	JS 86
	Harvest Index, %			
d1 : 20.000	43.63	53.29	18.13	19.71
d2 : 40.000	40.61	45.44	8.04	16.00
d3 : 80.000	36.49	43.80	6.80	11.37
d4 : 80.000 + Thinning	43.58	47.38	9.03	15.17
LSD (0.05)	3.48	---	2.99	4.25
LSD (0.01)	5.26	---	4.54	6.43
CV (%)	4.11	---	14.28	13.66

Table VII : Water use efficiency at Sidi El Aydi (SEA) and Jemaa Shaim (JS) in 1985 and 1986.

Treatment	Cropping Season and Location			
	SEA 85	JS 85	SEA 86	JS 86
	Kg Grain/cm Water Used			
d1 : 20.000	95.33	82.02	39.01	39.96
d2 : 40.000	110.63	110.88	20.46	39.50
d3 : 80.000	110.11	115.48	12.54	33.28
d4 : 80.000 + Thinning	132.75	107.46	19.47	32.11
LSD (0.05)	13.37	26.89	6.13	n.s.
LSD (0.01)	19.21	38.63	9.29	n.s.
CV (%)	7.45	16.17	13.43	16.50

ABSTRACT

The objective of this study was to determine if plant population reduction obtained either by seeding rate or by thinning plants at tasseling (practice of Moroccan farmers) can reduce the competition among plants during the reproductive and grain filling periods when moisture deficiency usually occurs. To reach this objectif experiments were conducted at Sidi El Aydi and Jemaa Shaim, Morocco, on alkaline vertisols under mean precipitation of 388 and 318 mm, respectively. Treatments consisted of four expected plant populations or densities: low -20,000 (d1), medium -40,000 (d2), high -80,000 (d3) and 80,000 plants / ha thinned at tasseling to 40,000 plants / ha (d4). Data collected were total dry matter; grain yield; yield components; midday leaf water potential; leaf diffusive resistance;leaf transpiration; and soil moisture. Increasing plant popuiation hastened soil water depletion, but total water used was not affected by population levels. Leaf diffusive resistance increased after silking. At the dent stage, the lowest leaf water potentials were -1.9, -2.2 and -3.3 MPa for the low, medium and high plant population, respectively. Critical value of leaf water potentiel for which leaves could not recover at night was -2.2 MPa. This value was never reached by the low plant population.

Drought during March and April probably hardened plants in 1985 and made them less sensitive to leaf water potential variation. Yield components, number of kernels/ear and seed weight tended to decrease with the increase of plant population. However, during a wet cropping season, high seed weight and number of kernels / ear obtained with the low plant population, were unable to compensate for the low plant population, consequently, yields remained low in this case. In a dry year, the two components mentioned above were relatively low, and yields were also very low. Thinning of plants at tasseling did not improve yields when the year was dry. In a relatively wet year thinnig improved yield (by 286 kg/ha) and its components.

From this study we can conclude tha thinning at tasseling is probably too late for saving water for the grain filling period and that 20,000 plants/ha in dry years and 40,000 plants/ha in wet years can be considered as optimum. Further investigation of plant populations is needed under these conditions

RESUME

L'objectif de cette étude est de déterminer si la diminution de la densité de peuplement, en jouant sur la dose de semis ou sur l'éclaircissage des plantes de maïs au stade panicule (pratique des agriculteurs marocains), peut réduire la compétition entre plantes au cours des périodes de reproduction et de remplissage de grain où le déficit hydrique est souvent observé.

Pour atteindre cet objectif, des essais sont conduits aux domaines expérimentaux de Sidi El Aydi et de Jamaa Shain (Maroc) sur des vertisols alcalins où les précipitations moyennes annuelles sont respectivement de 388 et 318 mm. Les densités de peuplement choisies pour cette étude sont : faible -20.000 (d1), intermédiaire -40.000 (d2), élevée -80.000 (d3) et 80.000 réduite à 40.000 pieds /ha (d4) par éclaircissage au stade panicule. Les données relevées sont, la matière sèche totale, le rendement grains, les principales composantes de rendement, le potentiel hydrique de la feuille, sa résistance stomatique, et sa transpiration ainsi que l'humidité du sol.

L'augmentation de la densité de peuplement a accélérée l'épuisement de l'eau du sol, mais la quantité d'eau totale utilisée n'a cependant pas été affectée par la variation de la densité de peuplement. Après floraison, la transpiration et le potentiel hydrique de la feuille ont chuté; cependant la résistance stomatique a augmenté au stade pateux. Les potentiels hydriques les plus bas sont respectivement -1.9, -2.2 et -3.3 MPa pour la faible, la moyenne et la plus forte densité de peuplement. La valeur critique du potentiel hydrique qui semble correspondre au point de flétrissement des feuilles est d'environ -2.2 MPa. Cette valeur n'est jamais atteinte par la faible densité de peuplement. Le manque de pluie en Mars et en Avril a probablement induit une certaine tolérance à la dessiccation chez le maïs en 1985 et a rendu les stomates des feuilles moins sensibles au variations du potentiel hydrique. Les composantes de rendement, nombre de grains/épi et le poids du grain ont tendance à chuter lorsque la densité de peuplement - pieds augmente. Cependant en année pluvieuse des niveaux élevés du poids du grain et du nombre de grains/épi chez la plus faible densité ne sont pas suffisants pour compenser le niveau bas du nombre de plantes/ha ; par conséquent les rendements dans ce cas sont restés faibles. En année sèche, les deux composantes mentionnées ci-dessus ainsi que le rendement grains sont généralement faibles surtout dans le cas de la plus forte densité de peuplement. De même l'éclaircissage au stade panicule ne semble pas améliorer les rendements. Néanmoins en année relativement plus arrosée, cet éclaircissage a amélioré le rendement grains (par 286 kg /ha) et ses composantes.

De cette étude, on peut conclure que la réduction du peuplement pieds par arrachage des plantes au stade panicule ne permet pas d'économiser de l'eau pour la période ultérieure du remplissage du grain et que les densités de 20.000 plantes/ha en année sèche et 40.000 en année pluvieuse peuvent être considérées comme optimales. Plus d'investigations sur les densités le peuplement sont nécessaires en zones semis-arides

ملخص

إن الهدف من هذه الدراسة هو معرفة ما إذا كان تخفيض كثافة نباتات الدرة عن طريق التقليل من كمية البذور أو التخفيف من عدد النباتات تؤدي إلى نقص التراحم بين النباتات خدمة التناصل وقت إمتلاء الحب.

ولتحقيق هذا الهدف، أجريت تجارب بمحطتي التجارب بسيدي العابدي بالشاوية وجمعة السحايم بعيدة، استعملت فيها أربع كثافات هي 20.000 (d1)، 40.000 (d2)، 80.000 (d3) و 80.000 في الهكتار التي تختلف في موعد التناصل من أجل الحصول على 40.000 نبتة في الهكتار (d4)، وقد أثبتت هذه التجارب خلال الموسمين الفلاحيين 1985-86 و 1986-87 حيث كان أحدهما (87-86) جافاً والأخر أكثر رطوبة.

وقد تم قياس المردود البيولوجي (المادة البياجة)، صردد المحبوب ومكوناته، جهد الماء الورقي، مناعة التفت، نفع النباتات ورطوبة التربة.

وتبين النتائج بأن الزيادة التدريجية في كثافة النباتات أسرع في طرح مشكل الماء، لكن الكمية المستهلكة من الماء لم تتأثر بتغيير الكثافة، ولوحظ أن نفاذ نفع النباتات بعد موعد التناصل وارتفاع جهد الماء الورقي ومناعة التفت عند مرحلة نضج البذور، في أواخر مرحلة إمتلاء المحب كان أعلى جهد الماء الورقي 3.1 ميكابامكمال بالنسبة لأقل كثافة، 2.26 بالنسبة للكثافة المتوسطة و 3.3 لأعلى كثافة، وقد لوحظت حالة الذبول عند الدرة حين كانت قيمة جهد الماء الورقي 2.2 ميكابامكمال تقاربها، وهذه قيمة لم تسجل أبداً في حالة الكثافة القليلة.

بالنسبة للسنة الجافة، وكانت نتائج الأسلار خلال شهري مارس وأبريل النبات من التكيف مع الحالة التي عرفتها آخر الدورة الرئاعية، أما مكونات المردود (عدد المحبوب في السنوية وذبذبة المحب) فقد سجلت تقريباً ملحوظاً حينما صارت الكثافة عالية، لكن خلال السنة المطرة، فإن كثرة المحب في السنوية نتيجة التقليل من كمية البذور (أقل كثافة) له توازن النقص في الكثافة، ولهذا بقى المردود متخفضاً، ويُمكن القول بأن تخفيف النباتات في مرحلة الإخصاب لم يحسن المردود خلال السنة الجافة عكس ما وقع خلال السنة المطرة.

الخلاصة من هذه الدراسة هي أن تخفيف الكثافة بالعملية اليدوية في مرحلة الإخصاب لا يمكن من احتصاد الماء وحرولة من أجل إستعماله لذوقت إمتلاء المحب، ولكن الإرداد بـ 20.000 نبتة في الهكتار في السنة الجافة و 40.000 نبتة في السنة المطرة.

REFERENCES BIBLIOGRAPHIQUES

- BABABOLA, O. AND C. OPUTA. 1981. Effects of planting patterns and population on water relations of maize. *J. Expl. Agric.* 17 : 97 - 104.
- BOYER, J.S. 1970. Differing sensitivity of photosynthesis to corn leaf water potentials in corn and soybean. *Plant Physiol.* 46 : 236-239.
- CLAASSEN, M.M. AND R.H. SHAW. 1970. Water deficits on corn. II. Grain components. *Agron. J.* 62 : 652-655.
- HARDEN, H.J., R.E. CARLSON, AND R.H. SHAW. 1982. Yield, yield components and nutrient content of corn grain as influenced by post-silking moisture stress. *Agron. J.* 74 : 275-278.
- PONELEIT, C.G. AND D.B. EGLI. 1979. Kernel growth and duration in maize as affected by plant density and genotype. *Crop Sci.* 19 : 385-388.
- TIMMONS, D.R; R.F. HOLT, AND J.T. MORAGHAN. 1966. Effect of corn population on yield, evapotranspiration and water use efficiency in the northwest cornbelt. *Agron. J.* 58 : 429-432.
- WALDREN, R.P. 1983. Corn. In : Teare, I.D. and M.M. Peet (Editors). *Crop Water Relations*, A wiley-Interscience Publ. N.Y., pp. 187-211.

BORDEREAU D'ENTREE DES DONNEES

AGRIS Formulaire 1(Rev. 5)F



001	C 1	VILLE	NUMERO DU NIVEAU	002	003	004	005
W 102	162			1 / 1		N C D	R N
TRN						NOUVEAU	SUSTITUE
TRANSLAT.	GENERIC					SUPRIME	
006	007	RN ou TRN de relation		MONOGRAPHIE	REPORT	ANALYTIQUE	REUNION
				NAME	ARTICLE	MONOGRAPHIE	DICTIO.
				DESIN	COLLECTIF	NAME	DOCTRINES
				FILM	REPORT	NAME	THESES
				CHTEAU	INFORM.	NAME	DISCUSSIONS
				CLASSE	ART. PUBL.	NAME	LEADERSHIP
				ENREGISTR.	BREVET	NAME	BIBLIOGRAPH.
				SERIE	PRT	NAME	CAFFES
				ART. SCIENT.		NAME	INDUSTRIE
						NAME	NON CONVENTION.
						NAME	SYNTHÈSE
008	F 61	(PRINCIPALE)	(SECONDAIRES)	CODE PAYS	TYPE BIBLIOGRAPHIQUE	NIVEAU BIBLIOGR.	INDICATEUR BIBLIOGRAPHIQUE
				(ENTREE REGIONALE)			

1 009 **A** Utiliser un bordereau pour chaque niveau bibliographique A, M ou C, cercle en 008, en partant du niveau le plus spécifique (c'est-à-dire la gauche) et reporter le code correspondant en 009. Pour le niveau bibliographique S, utiliser la section 2 du bordereau. Pour les descripteurs AOROVOC, les termes d'indexation du vocabulaire local et les résumés utiliser les sections 3 à 5 au verso.

	Entête	Données	(à dactylographier)
Auteur(s) Personne physique (Affiliation (s))	100	Karrou, M. (Aridoculture Centre, Settat (Maroc)) Minor, H.; Cray, L. (Univ. of Missouri, Columbia (Etats Unis). Dept. of Agronomy)	
Collectivité(s) auteur(s)	110		
Titre universitaire	111		
Titre anglais	Titre principal	Water use efficiency, growth and development response of maize to plant populations in semi-arid	
	Eléments secondaires	201 Morocco	
Réunion	Nom	210	
	Lieu	211	
	Date	213	
Titre original (Translit.)	Titre principal	230 Réponse de l'efficience d'utilisation de l'eau, de la croissance et de développement du maïs à la densité du peuplement au Maroc semi-aride	
	Eléments secondaires	231	
Edition (N°)	250		
No. Rapport/brevet	300		
Nos. accondaires	310		
ISBN/IPC	320		
Adresse bibliogra- phique	Lieu de publication	401	
	Editeur	402	
	Date de publication	403	
Collation	500		
Langue (s) du texte	600 (En)		
Notes	610 1601 Résumés (Ar, Fr, En) 7 tableaux, 4 fig., 7 ref.		
2 009 S	NIVEAU		

Titre de publication en arabe	Titre principal	230	Awamia (Maroc)
	Eléments secondaires	231	Revue Marocaine de la Recherche Agronomique
ISSN	320	ISSN 0522-2721	
Date de publication	403	(Jum 1992)	
Collation	500	(no 75) p. 3-17	
Notes	610		

009 9 / EN 009 0 / ES 009 9 / FR

Code de langue des descriptions (comme détaillées dans le champ ci-dessous)

Code de langage des termes d'indexation	Termes d'indexation	Définition (à encyclopédier)
009 9 / EN 800 / FR	ZEN MAIS; DENSITE DE SEMIS; BESOIN EN EAU; ECLAIRCISSAGE; STADE DE DEVELOPPEMENT VEGETAL; ZONE SEMI ARIDE; MAROC	(Séparer les descriptions par un point-virgule (;) et un espace. Faire précéder les propositions de nouveaux descripteurs par un point d'interrogation (?))
Autres descripteurs AGROVOC	/	(Laisser un espace après la barre oblique (/))

4 009 9 / Code de langue des termes d'indexation

Termes d'indexation du vocabulaire local	820

5 009 X / FR Code de langue du résumé

Langue du résumé en clair	850
Résumé	860 Etude ayant pour objectif de déterminer si la diminution de la densité de peuplement, en jouant sur la dose de semis ou sur l'éclaircissage des plantes de maïs au stade panicle, peut réduire la compétition entre plantes au cours des périodes de reproduction et de remplissage de grain où le déficit hydrique est souvent observé.

Etude ayant pour objectif de déterminer si la diminution de la densité de peuplement, en jouant sur la dose de semis ou sur l'éclaircissage des plantes de maïs au stade panicle, peut réduire la compétition entre plantes au cours des périodes de reproduction et de remplissage de grain où le déficit hydrique est souvent observé.

3113193

93-xx91

FIN

النهاية

19

مشاهد

VUES